

Stripe Charge Ordering in Triangular-Lattice Systems

Hiroaki Onishi and Takashi Hotta

Advanced Science Research Center, Japan Atomic Energy Research Institute, Tokai, Ibaraki 319-1195, Japan

(Dated: August 18, 2005)

We investigate the ground-state properties of a t_{2g} -orbital Hubbard model on a triangular lattice at electron density 5.5 by using numerical techniques. There appear several types of paramagnetic phases, but we observe in common that one or two orbitals among three orbitals become relevant due to the effect of orbital arrangement. It is found that charge stripes stabilized by the nearest-neighbor Coulomb interaction consist of antiferromagnetic/ferro-orbital chains for small Hund's coupling, while there occurs stripe charge ordering with ferromagnetic/antiferro-orbital chains for large Hund's coupling.

PACS numbers: 71.30.+h, 71.10.Fd, 75.30.Kz

Keywords: t_{2g} -orbital Hubbard model, triangular lattice, charge stripe

In recent years, a wide variety of exotic phenomena in layered cobalt oxides have attracted much attention in the research field of condensed-matter physics. In particular, the discovery of superconductivity in water-intercalated cobaltite $\text{Na}_{0.35}\text{CoO}_2 \cdot 1.3\text{H}_2\text{O}$ [1] has triggered current high activity on the study of superconductivity in frustrated triangular-lattice systems. On the other hand, magnetic transitions have also been observed in Na_xCoO_2 [2, 3]. In such materials, the competition between various types of ordered states should arise due to geometrical frustration. In addition, it has been stressed that the t_{2g} -orbital degree of freedom plays a crucial role in causing the rich phase diagram in the cobaltites. In fact, multiorbital models have been investigated to understand the mechanism of exotic superconductivity and novel magnetism in triangular-lattice cobalt oxides [4, 5, 6, 7]. To gain a deep insight into the behavior of such complex cobalt-based systems, it is important to clarify the characteristics of spin-charge-orbital states taking account of strong electron correlations

In this paper, we study the ground-state properties of the t_{2g} -orbital Hubbard model on a triangular lattice at electron density 5.5. The Hamiltonian is given by

$$\begin{aligned}
 H = & \sum_{\mathbf{i}, \mathbf{a}, \gamma, \gamma', \sigma} t_{\gamma\gamma'}^{\mathbf{a}} d_{\mathbf{i}\gamma\sigma}^\dagger d_{\mathbf{i}+\mathbf{a}\gamma'\sigma} \\
 & - (\Delta/3) \sum_{\mathbf{i}} (2\rho_{\mathbf{i}xy} - \rho_{\mathbf{i}yz} - \rho_{\mathbf{i}zx}) + V \sum_{\langle \mathbf{i}, \mathbf{j} \rangle} \rho_{\mathbf{i}} \rho_{\mathbf{j}} \\
 & + U \sum_{\mathbf{i}, \gamma} \rho_{\mathbf{i}\gamma\uparrow} \rho_{\mathbf{i}\gamma\downarrow} + (U'/2) \sum_{\mathbf{i}, \sigma, \sigma', \gamma \neq \gamma'} \rho_{\mathbf{i}\gamma\sigma} \rho_{\mathbf{i}\gamma'\sigma'} \\
 & + (J/2) \sum_{\mathbf{i}, \sigma, \sigma', \gamma \neq \gamma'} d_{\mathbf{i}\gamma\sigma}^\dagger d_{\mathbf{i}\gamma'\sigma'}^\dagger d_{\mathbf{i}\gamma\sigma'} d_{\mathbf{i}\gamma'\sigma} \\
 & + (J'/2) \sum_{\mathbf{i}, \sigma \neq \sigma', \gamma \neq \gamma'} d_{\mathbf{i}\gamma\sigma}^\dagger d_{\mathbf{i}\gamma'\sigma'}^\dagger d_{\mathbf{i}\gamma'\sigma'} d_{\mathbf{i}\gamma\sigma}, \quad (1)
 \end{aligned}$$

where $d_{\mathbf{i}\gamma\sigma}$ is the annihilation operator for an electron with spin σ in orbital γ ($=xy, yz, zx$) at site $\mathbf{i}$, $\rho_{\mathbf{i}\gamma\sigma} = d_{\mathbf{i}\gamma\sigma}^\dagger d_{\mathbf{i}\gamma\sigma}$, $\rho_{\mathbf{i}\gamma} = \sum_{\sigma} \rho_{\mathbf{i}\gamma\sigma}$, and $\rho_{\mathbf{i}} = \sum_{\gamma, \sigma} \rho_{\mathbf{i}\gamma\sigma}$. Note that the triangular lattice is arranged in the (x, y) plane (see




FIG. 1: (a) Triangular lattice with three directions depicted by arrows. An eight-site cluster enclosed by dotted lines is considered in our numerical calculations. (b) Schematic view of level splitting among t_{2g} orbitals.

Fig. 1). The hopping amplitudes are given by $t_{xy,xy}^{\mathbf{x}} = (dd\pi)$, $t_{yz,yz}^{\mathbf{x}} = t_{yz,zx}^{\mathbf{x}} = t_{zx,yz}^{\mathbf{x}} = 0$, $t_{zx,zx}^{\mathbf{x}} = (dd\pi)$ for the $\mathbf{x}$ direction, $t_{xy,xy}^{\mathbf{u}} = (9/16)(dd\sigma) + (1/4)(dd\pi)$, $t_{yz,yz}^{\mathbf{u}} = (3/4)(dd\pi)$, $t_{zx,zx}^{\mathbf{u}} = (1/4)(dd\pi)$, $t_{yz,zx}^{\mathbf{u}} = t_{zx,yz}^{\mathbf{u}} = (\sqrt{3}/4)(dd\pi)$ for the $\mathbf{u}$ direction, and $t_{\gamma,\gamma'}^{\mathbf{v}} = t_{\gamma,\gamma'}^{\mathbf{u}}$ except for $t_{yz,zx}^{\mathbf{v}} = t_{zx,yz}^{\mathbf{v}} = (-\sqrt{3}/4)(dd\pi)$ for the $\mathbf{v}$ direction. Note that there is no hopping between the $a_{1g}(xy)$ and $e'_g(yz \text{ and } zx)$ orbitals due to their different symmetry. Hereafter, we take $(dd\pi)=1$ as the energy unit. Δ is the level splitting between the a_{1g} and e'_g orbitals, and V is the nearest-neighbor Coulomb interaction. In the local interactions, U , U' , J , and J' are intraorbital Coulomb, interorbital Coulomb, exchange, and pair hopping interactions, respectively. Note the well-known relations $J=J'$ and $U=U'+J+J'$. We analyze the model for eight-site ($N=8$) systems with periodic boundary conditions by the Lanczos diagonalization. In this paper, we set $(dd\sigma)=2$, $\Delta=6$, and $U'=20$, and investigate the dependence on J and V . The dependence on $(dd\sigma)$, Δ , and U' will be discussed elsewhere in future.

Our result is summarized in Fig. 2(a), which is the ground-state phase diagram in the (V, J) plane. Comparing the lowest energies in the subspaces of the z component of the total spin, we determine the total spin in the ground state. Then, we find that for a large set of parameters, the ground state is a paramagnetic (PM) phase characterized by total spin zero, but there appear several types of PM phases in terms of spin-charge-orbital configurations. Among them, the phase I is characterized by a stripe, in which ferromagnetic (FM) spin chains are




FIG. 2: (a) Ground-state phase diagram in the (V, J) plane for $\Delta=6$ and $U'=20$ at electron density 5.5. Inset denotes schematic view of the charge stripe and spin configuration in each phase. The size of circle indicates the degree of the charge density. (b) Electron densities in the xy , yz , and zx orbitals. (c) Electron density in the a orbital, represented by the linear combination of the yz and zx orbitals.

coupled with each other in an antiferromagnetic (AFM) manner, as shown in the inset of phase I. Note that the charge is uniformly distributed in the background of ferro-orbital (FO) arrangement.

On the other hand, the phases II and III exhibit charge order. To clarify what types of charge configurations emerge in the present system, we investigate the charge structure factor. When we examine the V dependence of the charge structure factor for $J=0$, it is clearly observed that it develops abruptly around at $V=10$, indicating the stabilization of a specific charge structure. The results suggest a charge stripe composed of one-dimensional chains, as shown in the inset of phase II. Namely, the spin frustration is removed due to charge ordering. It should be noted that a similar stripe charge pattern has been suggested for a CoO_2 triangular-lattice model by Hartree-Fock calculations [7]. Note that in the phase II, two of three orbitals are found to be fully occupied due to the effect of FO arrangement, similar to the phase I.

Now we consider the effect of J on the spin structure in the stripe charge state. For this purpose, it is useful to measure the spin structure factor. For $V=12$, we observe a transition in the spin structure factor at around $J=0.6$, but the stripe charge configuration is not changed. For $J<0.6$, we find AFM correlations in each one-dimensional charge stripe, as shown in the inset of phase II. On the other hand, for $J>0.6$, the spin configuration changes

to a FM one, as shown in the inset of phase III. Weak antiferro-orbital (AFO) correlations exist in the chains, since the electrons can gain kinetic energy due to the so-called double-exchange mechanism.

In order to identify the orbital arrangement, it is convenient to evaluate the electron densities, $n_\gamma = \sum_i \langle \rho_{i\gamma} \rangle / N$, in the xy , yz , and zx orbitals. In Fig. 2(b), we show n_γ vs. J for $V=12$. It is observed that the lower xy orbital is fully occupied irrespective of J , while the electron densities in the yz and zx orbitals change suddenly at around $J=0.6$. For $J<0.6$, electrons favorably occupy the zx orbital, but the zx orbital is not fully occupied. On the other hand, the yz and zx orbitals are equally occupied for $J>0.6$.

However, in order to determine the orbital structure, we need to pay due attention to the definition of orbitals. Here, by using an angle θ_i to characterize the orbital shape, we introduce operators for a and b orbitals such as $d_{ia\sigma} = e^{i\theta_i/2} [\cos(\theta_i/2) d_{i,yz,\sigma} + \sin(\theta_i/2) d_{i,zx,\sigma}]$ and $d_{ib\sigma} = e^{i\theta_i/2} [-\sin(\theta_i/2) d_{i,yz,\sigma} + \cos(\theta_i/2) d_{i,zx,\sigma}]$. In Fig. 2(c), we show the electron density in the a orbital with $\theta_i = \theta$. We find that for $J=0$, the a orbital of $\theta=4\pi/3$ is fully occupied. Namely, only one orbital becomes relevant for small J . This FO arrangement is also found in the phase I. On the other hand, for $J=1$, the electron density does not change even when θ is varied, indicating that two orbitals remain active.

Note that we have suggested the stripe charge order from the 8-site calculations, but the shape of the stripe may change from straight to zigzag. To confirm this point, it is necessary to perform the calculations in larger-sized systems, for instance, using a model including only e'_g orbitals. This is among possible future developments.

In summary, in the PM phase, one or two orbitals among the t_{2g} orbitals become relevant due to the effect of orbital arrangement. The charge stripe stabilized by V consists of AFM/FO chains for small J , while it changes to FM/AFO ones with increasing J .

We thank E. Dagotto for discussions. T.H. is supported by the Japan Society for the Promotion of Science and by the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

-
- [1] K. Takada *et al.*, Nature (London) **422**, 53 (2003).
 - [2] T. Motohashi *et al.*, Phys. Rev. B **67**, 064406 (2003).
 - [3] J. Sugiyama *et al.*, Phys. Rev. B **69**, 214423 (2004).
 - [4] M. Mochizuki *et al.*, Phys. Rev. Lett. **94**, 147005 (2005).
 - [5] W. Koshibae and S. Maekawa, Phys. Rev. Lett. **91**, 257003 (2003).
 - [6] M. Indergand *et al.*, Phys. Rev. B **71**, 214414 (2005).
 - [7] T. Mizokawa, New J. Phys. **6**, 169 (2004).